



# MALARIA RISK AREAS



## North, Central & South America

- Argentina 22
- Bahamas 9
- Belize 3
- Bolivia 20
- Brazil 18
- Colombia 13
- Costa Rica 7
- Dominican Republic 11
- El Salvador 4
- Ecuador 12
- French Guiana 17
- Guatemala 2
- Guyana 15
- Haiti 10
- Honduras 5
- Mexico 1
- Nicaragua 6
- Panama 8
- Paraguay 21
- Peru 19
- Suriname 16
- Venezuela 14

## Africa

- Algeria 25
- Angola 61
- Banin 47
- Botswana 72
- Burkina Faso 44
- Burundi 57
- Cameroon 49
- Cape Verde 36
- Cent. African Rep. 50
- Chad 30
- Comoros 65
- Republic of Congo 54
- Democratic Republic of Congo 55
- Djibouti 33
- Egypt 26
- Equatorial Guinea 51
- Eritrea 32
- Ethiopia 34
- Gabon 53
- Gambia 38
- Ghana 45
- Guinea 40

- Guinea-Bissau 39
- Ivory Coast 43
- Kenya 59
- Liberia 42
- Libya 25
- Madagascar 69
- Malawi 63
- Mali 28
- Mauritania 27
- Mauritius 67
- Morocco 23
- Mozambique 64
- Namibia 71
- Niger 29
- Nigeria 48
- Reunion 68
- Rwanda 56
- Sao Tome & Principe 52
- Senegal 37
- Seychelles 66
- Sierra Leone 41
- Somalia 35

- South Africa 73
- Sudan 31
- Swaziland 74
- Tanzania 60
- Togo 46
- Uganda 58
- Zambia 62
- Zanzibar 107
- Zimbabwe 70

## Asia & Oceania


- Afghanistan 84
- Azerbaijan 76
- Bangladesh 90
- Bhutan 91
- Cambodia 99
- China 93
- Hong Kong 94
- India 86
- Indonesia 102
- Iran 83
- Iraq 78
- Laos 97
- Malaysia 100
- Maldives 87
- Myanmar (Burma) 92
- Nepal 89
- Oman 81
- Pakistan 85
- Papua New Guinea 104
- Philippines 103
- Saudi Arabia 79
- Singapore 101
- Solomon Islands 105
- Sri Lanka 88
- Syria 77
- Taiwan 95
- Thailand 98
- Turkey 75
- United Arab Emirates 82
- Vanuatu 106
- Vietnam 96
- Yemen 80

### MALARIA RISK MAP FOR SOUTH AFRICA

To significantly reduce your risk, take precautionary measures against mosquito bites throughout the year in ALL RISK areas.

Where malaria chemoprophylaxis is indicated, mefloquine or atovaquone-proguanil or doxycycline should be used.

- Low Risk**  
Only non-drug measures to prevent mosquito bites are recommended from September to May
- Moderate Risk**  
Antimalarial drugs are recommended from September to May for all travellers
- Malaria risk in neighbouring countries**  
For further information, please consult the WHO travel health guidelines at [www.who.int/ith/en/](http://www.who.int/ith/en/)



Please contact our Head Medical Nurse for more information:  
021 507 6355